

Rosa Cid-Muñoz · Manuel Pedreira

Classification of Incidence Scrolls(I)

Abstract. The aim of this paper is to obtain a classification of scrolls of genus 0 and 1, which are defined by a one-dimensional family of lines meeting a certain set of linear spaces in $\mathbf{P}^n$. These ruled surfaces will be called incidence scrolls and such a set will be the base of the incidence scroll. Unless otherwise stated, we assume that the base spaces are in general position.

Introduction

Throughout this paper, the base field for algebraic varieties is $\mathbb{C}$. Let $\mathbf{P}^n$ be the n -dimensional complex projective space and $G(l, n)$ the Grassmannian of l -planes in $\mathbf{P}^n$. Then $R_g^d \subset \mathbf{P}^n$ denotes a scroll of degree d and genus g . We will follow the notation of [3].

Projective ruled surfaces with the property that the generators meet a suitable number of linear spaces in $\mathbf{P}^n$ are known classically. The first examples of this situation are the smooth quadric surface in $\mathbf{P}^3$ and the elliptic quintic scroll in $\mathbf{P}^4$ formed by the lines which meet five planes in general position. Although this is not so in all the ruled surfaces, there does not exist a classification of such ruled surfaces. Our purpose is to investigate the scrolls in $\mathbf{P}^n$ whose generators meet a certain set of linear spaces in general position. The ruled surfaces so obtained will be called incidence scrolls, and such a set will be a base of the incidence scroll.

It is useful to represent a scroll in $\mathbf{P}^n$ by a curve $C \subset G(1, n) \subset \mathbf{P}^N$. The lines which intersect a given projective space $\mathbf{P}^r$ in $\mathbf{P}^n$ are represented by the points of the special

Mathematics Subject Classifications: 14J26 (14H25, 14H45)

Authors' address: Departamento de Algebra, Universidad de Santiago de Compostela, 15782 Santiago de Compostela, Galicia, Spain. Phone: 34-81563100-ext.13152. Fax: 34-81597054.e-mail: rosacid@usc.es; pedreira@zmat.usc.es

Schubert variety $\Omega(\mathbf{P}^r, \mathbf{P}^n)$. Each $\Omega(\mathbf{P}^r, \mathbf{P}^n)$ is the intersection of $G(1, n)$ with a certain linear space in $\mathbf{P}^N$. Since $G(1, n)$ has dimension $2n - 2$ and we need a curve, we must impose $2n - 3$ linear conditions on $G(1, n)$. Consequently, the choice of linear spaces is not arbitrary. Any set of linear spaces in $\mathbf{P}^n$ which imposes $2n - 3$ linear conditions on $G(1, n)$ is the base of a certain incidence scroll. The background about Schubert varieties can be found in [6].

We will expose a method to know when a scroll is determined by incidences. The affirmative solution would allow us to obtain a base for each incidence scroll (there is really only one way to choose this base). This is possible because the families of directrix curves provide a natural and intrinsic characterization of the incidence scrolls.

Our first step will be to give some general properties of ruled surfaces. For more details we refer the reader to [3]. We do it in section 1, where we define a particular class of surfaces, the scrolls. In section 2, having introduced the notion of incidence scroll, we have compiled some basic properties of such a scroll. The degree of the scroll given by a general base is provided by Giambelli's formula. Our main contribution is to establish (IS) which together with 2.5 - 2.7 gives the first base spaces of the incidence scroll.

Section 3 is devoted to the study of deformations of a given incidence scroll. These form a powerful tool for obtain a finite family of incidence scrolls from a given one. If the incidence scroll $R_g^d \subset \mathbf{P}^n$ breaks up into $R_{g_1}^{d_1} \subset \mathbf{P}^r$ and $R_{g_2}^{d_2} \subset \mathbf{P}^s$ with δ generators in common, then $d = d_1 + d_2$ and $g = g_1 + g_2 + \delta - 1$.

Sections 4 and 5 contain our main results about rational and elliptic incidence scrolls. In the former section, we show that the only rational scrolls determined by incidences are those of general type and those with a directrix line. In the latter section, we give all the elliptic incidence scrolls. In each case, we find the base of the scroll. Accordingly, we will apply the following results to obtain a large number of incidence scrolls:

- Bertini's theorem: if Λ is a reduced linear system on a complete nonsingular variety V such that $\dim(\Phi(V)) \geq 2$, then the generic element of Λ is irreducible (see [4], p. 252).
- A family of closed subschemes $f: X \subset T \times \mathbf{P}^n \longrightarrow T$ is flat if and only if the Hilbert polynomials of the fibres are the same and hence also the arithmetic genus. Every algebraic family of normal varieties parametrized by a nonsingular curve T over an algebraically closed field is a flat family of schemes. For any flat family, having a point $t \in T$ with X_t birationally equivalent to X_0 , we say that X is a (global) deformation of X_0 (see [3], pp. 256-266).

We then prove the main theorem of this sections, which determines all the incidence scrolls of genus 0 and 1.

Theorem(see Theorems 4.2 and 5.1) *Let X be a ruled surface over the curve C of genus $g = 0, 1$. Let $H \sim C_o + \mathfrak{b}f$ be a very ample divisor on X with $m = \deg(\mathfrak{b})$ and let $\Phi_H : X \rightarrow R_g^{2m-e} \subset \mathbf{P}^{2m-e-2g+1}$ be the closed immersion defined by $|H|$. Then R_g^{2m-e} is an incidence scroll if and only if it satisfies one of the following conditions:*

1. $g = 0$ and $e = 0, 1$;
2. $g = 0$ and $m = e + 1$;
3. $g = 1$, $e = -1$ and $m = 2$;
4. $g = 1$, $\mathfrak{e} \sim 0$ and $m = 4$;
5. X decomposable, $g = 1$, $0 \leq e \leq 3$ and $m = e + 3$.

It is to be expected that the treatment of general case will be answered shortly. A paper on incidence scrolls in $\mathbf{P}^n$ is to be written. The results on this paper belong to the Ph.D. thesis of the first author whose advisor is the second one.

We thank the referee for many detailed remarks of a first version of this paper.

1. Ruled Surfaces

A ruled surface is a surface X together with a surjective morphism $\pi: X \longrightarrow C$ to a (connected nonsingular) curve C such that the fibre X_y is isomorphic to $\mathbf{P}^1$ for every point $y \in C$, and such that π admits a section. There exists a locally free sheaf $\mathcal{E}$ of rank 2 on C such that $X \cong \mathbf{P}(\mathcal{E})$ over C . Conversely, every such $\mathbf{P}(\mathcal{E})$ is a ruled surface over C . Let $\mathcal{O}_X(1)$ be the invertible sheaf $\mathcal{O}_{\mathbf{P}(\mathcal{E})}(1)$. Then there is an one-to-one correspondence between sections $\sigma: C \longrightarrow X$ and surjections $\mathcal{E} \longrightarrow \mathcal{L} \longrightarrow 0$, where $\mathcal{L}$ is an invertible sheaf on C , given by $\sigma^*(\mathcal{O}_X(1))$. Moreover, if D is any section of X , corresponding to a surjection $\mathcal{E} \longrightarrow \mathcal{L} \longrightarrow 0$, and if $\mathcal{L} = \mathcal{L}(\mathfrak{d})$ for some divisor $\mathfrak{d}$ on C , then $\deg(\mathfrak{d}) = C_o \cdot D$, and D and $C_o + (\mathfrak{d} - \mathfrak{e})f$ are linearly equivalent, written $D \sim C_o + (\mathfrak{d} - \mathfrak{e})f$.

It is possible to write $X \cong \mathbf{P}(\mathcal{E})$ where $\mathcal{E}$ is a locally free sheaf on C with the property that $H^0(\mathcal{E}) \neq 0$ but for all invertible sheaves $\mathcal{L}$ on C with $\deg \mathcal{L} < 0$, we have $H^0(\mathcal{E} \otimes \mathcal{L}) = 0$. In this case we say $\mathcal{E}$ is normalized. Let $\mathfrak{e}$ be the divisor on C corresponding to the invertible $\bigwedge^2 \mathcal{E}$ and let $e = -\deg \mathfrak{e}$. Furthermore there is a section $\sigma_0: C \longrightarrow X$ with image C_o such that $\mathcal{L}(C_o) \cong \mathcal{O}_X(1)$. Fix such a section C_o of X . If $\mathfrak{b}$ is any divisor on C , then we denote the divisor $\pi^*\mathfrak{b}$ on X by $\mathfrak{b}f$, by abuse of notation. Thus any element of $\text{Pic}(X)$ can be written $aC_o + \mathfrak{b}f$ with $a \in \mathbb{Z}$ and $\mathfrak{b} \in \text{Pic}(C)$.

Let X be a ruled surface over the curve C of genus g , determined by a normalized locally free sheaf $\mathcal{E}$. If $\mathcal{E}$ is decomposable, then $\mathcal{E} \cong \mathcal{O}_C \oplus \mathcal{L}$ for some $\mathcal{L}$ with $\deg \mathcal{L} \leq 0$. All values of $e \geq 0$ are possible. If $\mathcal{E}$ is indecomposable, then $-g \leq e \leq 2g - 2$.

A scroll is a ruled surface embedded in $\mathbf{P}^N$ in such a way that the fibres f have degree 1. If we take a very ample divisor on X , $D \sim aC_o + \mathfrak{b}f$, then the embedding $\Phi_D: X \longrightarrow R_g^d \subset \mathbf{P}^N = \mathbf{P}(H^0(\mathcal{O}_X(D)))$ determines a scroll when $D \cdot f = a = 1$. Throughout this work, we will restrict our attention to very ample divisors so that $\Phi_D(X)$ be a scroll.

2. Definition of Incidence Scroll

Definition 2.1 A scroll $R \subset \mathbf{P}^n$ is said to be an *incidence scroll* if R is generated by the lines which meet a certain set $\mathcal{B}$ of linear spaces in $\mathbf{P}^n$. Such a set is called a *base* of the incidence scroll.

Since a scroll $R_g^d \subset \mathbf{P}^n$ is represented by a curve $C_g^d \subset G(1, n)$, there is another definition of incidence scroll which is equivalent to the previous one.

Definition 2.2 A scroll $R \subset \mathbf{P}^n$ is an *incidence scroll* if the correspondent curve in $G(1, n)$ is an intersection of special Schubert varieties $\Omega(\mathbf{P}^r, \mathbf{P}^n)$, $0 \leq r < n - 1$.

Such a base will be denoted by:

$$\mathcal{B} = \{\mathbf{P}^{n_1}, \mathbf{P}^{n_2}, \dots, \mathbf{P}^{n_r}\}$$

or we will write it simply $\mathcal{B}$ when no confusion can arise, where $n_1 \leq n_2 \leq \dots \leq n_r$, i.e., $C = \bigcap_{i=1}^r (\Omega(\mathbf{P}^{n_i}, \mathbf{P}^n)) \subset G(1, n)$. If $n_1 = \dots = n_i$, $n_{i+1} = \dots = n_j$, ... and $n_{j+1+k} = \dots = n_r$, then we write it $\mathcal{B} = \{i\mathbf{P}^{n_1}, (j-i)\mathbf{P}^{n_j}, \dots, (r-j-k)\mathbf{P}^{n_r}\}$, for short.

We can certainly assume that $n \geq 3$. This involves no loss of generality because the only scroll in $\mathbf{P}^2$ is $\mathbf{P}^2$. The plane can be obtained as incidence scroll in $\mathbf{P}^n$ if we take $\mathcal{B} = \{\mathbf{P}^0, (n-2)\mathbf{P}^{n-2}\}$. If we want to see it as nondegenerate surface, we must work in $\mathbf{P}^2$ with $\mathcal{B} = \{\mathbf{P}^0\}$ (i.e., $\mathbf{P}^1 = \Omega(\mathbf{P}^0, \mathbf{P}^2)$).

We are interested in incidence scrolls where the base is formed by linear spaces in general position. By general position we will mean that $(\mathbf{P}^{n_1}, \dots, \mathbf{P}^{n_r}) \in \mathcal{W} = G(n_1, n) \times \dots \times G(n_r, n)$ is contained in a nonempty open subset of $\mathcal{W}$. Therefore, unless otherwise stated, we will work with general base spaces. For simplicity of notation, we abbreviate it to base in general position.

Theorem 2.3 The intersection $\bigcap_{j=1}^m \Omega(\mathbf{P}^{n_j}, \mathbf{P}^n) \subset G(l, n)$ of the special Schubert varieties associated to linear subspaces $\mathbf{P}^{n_j}$, $j = 1, \dots, m$, of dimension n_j such that $(l +$

$1)(n-l) - \sum_{j=1}^m (n-n_j-1) > 0$, is connected. Moreover, the intersection is irreducible of dimension $(l+1)(n-l) - \sum_{j=1}^m (n-n_j-1)$ for a general choice of the subspaces $\mathbf{P}^{n_j}$.

Proof. See [5], Theorem 1.1.

With the previous theorem, we can prove immediately an important consequence which provides a characterization of incidence scrolls.

Proposition 2.4 *The intersection $C = \bigcap_{i=1}^r \Omega(\mathbf{P}^{n_i}, \mathbf{P}^n)$ of special Schubert varieties associated to linear spaces $\mathbf{P}^{n_i}$, $i = 1, \dots, r$ in general position is a conex irreducible curve of $G(1, n)$ if and only if it verifies the following equality*

$$rn - (n_1 + n_2 + \dots + n_r) - r = 2n - 3 \quad (IS)$$

Proof. " $\Rightarrow$ " Since $C \subset G(1, n)$ is a conex irreducible curve, we have $1 = 2(n-1) - rn + r + \sum_{i=1}^r n_i$ (see Theorem 2.3) $\iff rn - \sum_{i=1}^r n_i - r = 2n - 3$.

" $\Leftarrow$ " Since $2(n-1) - \sum_{i=1}^r n - n_i - 1 > 0$, we conclude that C is connected, by Theorem 2.3. Moreover, we easily see from Theorem 2.3 that it is irreducible of dimension 1 for a general choice of the $\mathbf{P}^{n_i}$, $i = 1, \dots, r$. ■

Each $\Omega(\mathbf{P}^{n_i}, \mathbf{P}^n)$ is the intersection of $G(1, n)$ with a linear space in $\mathbf{P}^N$. Moreover, such a linear space is a hyperplane if and only if we have $n_i = n - 2$. If every dimension of the base spaces is equal to $n - 2$, then we will need exactly $(2n - 3) \mathbf{P}^{n-2}$'s for obtain a curve in $G(1, n)$.

Some relevant properties of incidence scrolls will be indicated. These are elementary but of great importance if we want to develop a theory about incidence scrolls.

1. A hyperplane does not impose conditions on $G(1, n)$ because $\Omega(\mathbf{P}^{n-1}, \mathbf{P}^n) = G(1, n)$.

Then we can assume $n_r \leq n - 2$.

2. If $n_i + n_j < n - 1$ then the scroll is degenerated. Under the above assumptions,

$$\Omega(\mathbf{P}^{n_i}, \mathbf{P}^n) \cap \Omega(\mathbf{P}^{n_j}, \mathbf{P}^n) = \Omega(\mathbf{P}^{n_i}, \mathbf{P}^{n_i+n_j+1}) \cap \Omega(\mathbf{P}^{n_j}, \mathbf{P}^{n_i+n_j+1}).$$

3. A nondegenerate (irreducible) incidence scroll cannot have double points. In other case, if P is a double point, there are two generators of the scroll g_1, g_2 through P . Thus, the incidence scroll is not irreducible, containing the plane pencil determined by them. In $G(1, n)$, we have $\bigcap_{i=1}^r \Omega(\mathbf{P}^{n_i}, \mathbf{P}^n) = \Omega(P, \mathbf{P}^2) \cup (\bigcap_{j=1}^s \Omega(\mathbf{P}^{m_j}, \mathbf{P}^m))$.
4. The generators of the scroll meet each base space in points of a directrix curve.
5. The degree of the scroll can be calculated by a one-to-one correspondence between any two of the directrix curves. It can also be computed by Giambelli's formula, i.e., it is the number of lines in $\mathbf{P}^n$ which intersect $\mathbf{P}^{n_1}, \mathbf{P}^{n_2}, \dots, \mathbf{P}^{n_r}$ and a generic $\mathbf{P}^{n-2}$. If $\deg(R) = d$, then we have the following equality of Schubert cycles: $\Omega(n_1, n) \cdots \Omega(n_r, n) = d\Omega(0, 2)$.

From now on, we will talk about a decomposable incidence scroll if the corresponding ruled surface $X = \mathbf{P}(\mathcal{E})$ has $\mathcal{E}$ decomposable. If $\mathcal{E}$ is indecomposable, then we will talk of an indecomposable incidence scroll.

Let $X = \mathbf{P}(\mathcal{E})$ be a ruled surface over the curve C of genus g , determined by a decomposable normalized bundle $\mathcal{E} \cong \mathcal{O}_C \oplus \mathcal{O}_C(\mathbf{e})$ such that $\deg(-\mathbf{e}) = e \geq 0$. Let $H \sim C_o + \mathbf{b}f$ be the very ample divisor on X with $m = \deg(\mathbf{b})$ which gives the immersion of the ruled surface as the scroll $R_g^d \subset \mathbf{P}^n$ such that $d = 2m - e$ and $n = 2(m - g) - e + 1 + i$, being i the speciality of the scroll ($i = h^1(\mathcal{O}_C(\mathbf{b})) + h^1(\mathcal{O}_C(\mathbf{b} + \mathbf{e}))$). Geometrically, X has two disjoint directrix, denoted by C_o and C_1 , such that $C_1 \sim C_o - \mathbf{e}f$. Moreover, it is easy to check that these satisfy $\phi_{\mathbf{b}+\mathbf{e}}: C_o \longrightarrow C_g^{m-e} \subset \mathbf{P}^{m-e-g+i_1}$ and $\phi_{\mathbf{b}}: C_1 \longrightarrow C_g^m \subset \mathbf{P}^{m-g+i_2}$, being $i_1 = h^1(\mathcal{O}_C(\mathbf{b} + \mathbf{e}))$ and $i_2 = h^1(\mathcal{O}_C(\mathbf{b}))$. Therefore $i = i_1 + i_2$ and $\mathbf{P}^{m-e-g+i_1} \cap \mathbf{P}^{m-g+i_2} = \emptyset$.

Proposition 2.5 *If $R_g^d \subset \mathbf{P}^n$ is a decomposable incidence scroll with base in general position, then $\mathbf{P}^{m-e-g+i_1}$ and $\mathbf{P}^{m-g+i_2}$ are base spaces.*

Proof. Suppose $\mathcal{B}$ a base with $\sum_{i=1}^r (n - n_i - 1) = 2n - 3$ such that $\mathbf{P}^{n_i}$'s, $i = 1 \dots, r$, are in general position and $m - e - g + i_1 < n_i \leq n - 2$. Since the scroll has a minimum directrix curve $C_g^{m-e} \subset \mathbf{P}^{m-e-g+i_1}$, we can take a generic hyperplane $\mathbf{P}^{n-1}$ through $\mathbf{P}^{m-e-g+i_1}$. Then there are m lines in $\mathbf{P}^{n-1}$ which meet $\mathbf{P}^{n_1-1}, \mathbf{P}^{n_2-1}, \dots$ and $\mathbf{P}^{n_r-1}$. Since each $\mathbf{P}^{n_i-1}$ imposes $n - n_i - 1$ independent conditions on $\mathbf{P}^{n-1}$, it is impossible.

$\mathbf{P}^{m-e-g+i_1}$ imposes $m - g + i_2$ conditions on $G(1, n)$. Then we can consider more subspaces to form the base. There is a $\mathbf{P}^{m-g+i_2} \in \mathcal{B}$. For otherwise, suppose $\mathcal{B} = \{\mathbf{P}^{m-e-g+i_1}, \mathbf{P}^{n_1}, \dots, \mathbf{P}^{n_r}\}$ (in general position) with $m - g + i_2 < n_i \leq n - 2$. Then we can take a generic hyperplane $\mathbf{P}^{n-1}$ such that $\mathbf{P}^{m-g+i_2} \subset \mathbf{P}^{n-1}$ and arguments similar to the above imply that $\mathcal{B}$ is not the base of $R_g^d \subset \mathbf{P}^n$. ■

Theorem 2.6 *Let $R_g^d \subset \mathbf{P}^n$ be an incidence scroll with base in general position. Then:*

$$R \text{ is decomposable} \iff \mathbf{P}^{n_1} \cap \mathbf{P}^{n_2} = \emptyset.$$

Proof. " $\Leftarrow$ " Since $\mathbf{P}^{n_1} \cap \mathbf{P}^{n_2} = \emptyset$, the directrix curves $C_g^{d_1} \subset \mathbf{P}^{n_1}$ and $C_g^{d_2} \subset \mathbf{P}^{n_2}$ satisfy $C_g^{d_1} \cap C_g^{d_2} = \emptyset$, i.e., R_g^d is decomposable.

" $\Rightarrow$ " By Proposition 2.5, there are two directrix curves $C_g^{d_1} \subset \mathbf{P}^{n_1}$ and $C_g^{d_2} \subset \mathbf{P}^{n_2}$ such that $C_g^{d_1} \cap C_g^{d_2} = \emptyset$. Suppose $n_j = d_j - g + i_j$ for $j = 1, 2$. Since $n = n_1 + n_2 + 1$ and $\mathbf{P}^{n_1}$ and $\mathbf{P}^{n_2}$ are in general position, $\mathbf{P}^{n_1} \cap \mathbf{P}^{n_2} = \emptyset$. ■

If R is decomposable, $\mathbf{P}^{m-e-g+i_1}$ and $\mathbf{P}^{m-g+i_2}$ impose $2(m - g) - e + i$ conditions on $G(1, n)$. Since $2n - 3 - (2(m - g) - e + i) = n - 2 \geq 1$, we must consider more subspaces to form the base.

In particular, if $e \leq 2g - 2$, then $h^0(\mathcal{O}_X(C_o - \mathfrak{e}f)) = e + 2 - g$. Moreover, if $g = 0, 1$, then we can show (by Bertini's theorem) that there are $e + 2 - g$ linearly independent directrix curves $C \sim C_o - \mathfrak{e}f$, which generates $|C_o - \mathfrak{e}f|$.

Proposition 2.7 *Let $R_g^{2m-e} \subset \mathbf{P}^{2(m-g)-e+1+i}$ be a decomposable incidence scroll with base in general position.*

- (a) If $\mathfrak{e} \not\sim 0$, then there are $(e + 2 - g) \mathbf{P}^{m-g+i_2}$'s in $\mathcal{B}$, whenever possible, i.e., when $m - g + i_2 + (e + 2 - g)(m - e - g + i_1) \leq 4(m - g) - 2e + 2i - 1$ and the generic curve in $|C_o - \mathfrak{e}f|$ is irreducible.
- (b) If $\mathfrak{e} \sim 0$, then there are $3 \mathbf{P}^{m-e-g+i_1}$'s in $\mathcal{B}$.

Proof. (a) We proceed by induction in $j = e + 2 - g$. The proposition is true for $j = 1$, by Proposition 2.5. Supposing the proposition true for j , we prove it for $(j + 1) \mathbf{P}^{m-g+i_2}$'s. Let $\mathcal{B} = \{\mathbf{P}^{m-e-g+i_1}, \mathbf{P}^{n_1}, \mathbf{P}^{n_2}, \dots, \mathbf{P}^{n_r}\}$ be a base of the scroll in general position with $m - g + i_2 = n_1 = \dots = n_j < n_{j+1} \leq \dots \leq n_r \leq 2(m - g) - e + i - 1$. Since the scroll has at least one directrix curve $C_g^m \subset \mathbf{P}^{m-g+i_2} \notin \mathcal{B}$, which is linearly independent from the other directrix curves contained in base spaces of dimension $m - g + i_2$, we can take a generic hyperplane $\mathbf{P}^{2(m-g)-e+i}$ through $\mathbf{P}^{m-g+i_2}$. Then there are $m - e$ lines in $\mathbf{P}^{2(m-g)-e+i-j}$ which meet $\mathbf{P}^{m-e-g+i_1-1}, \mathbf{P}^{n_1-j}, \dots, \mathbf{P}^{n_j-j}, \mathbf{P}^{n_{j+1}-j-1}, \dots$ and $\mathbf{P}^{n_r-j-1}$, which is impossible.

The proof for (b) is similar. ■

3. Degeneration

Proposition 3.1 *Let $R_g^d \subset \mathbf{P}^n$ be an incidence scroll with base $\mathcal{B}$ in general position. Suppose that $\mathbf{P}^{n_i}$ and $\mathbf{P}^{n_j}$ lie in a hyperplane $\mathbf{P}^{n-1}$ and have in common $\mathbf{P}^m$, $m = n_i + n_j - n + 1$. Then the scroll breaks up into:*

- $R_{g_1}^{d_1} \subset \mathbf{P}^n$ with base $\dot{\mathcal{B}} = \{\mathbf{P}^m, \mathbf{P}^{n_1}, \dots, \widehat{\mathbf{P}^{n_i}}, \dots, \widehat{\mathbf{P}^{n_j}}, \dots, \mathbf{P}^{n_r}\}$ (which is possibly degenerate);
- $R_{g_2}^{d_2} \subset \mathbf{P}^{n-1}$ with base $\ddot{\mathcal{B}} = \{\dots, \mathbf{P}^{n_{i-1}-1}, \mathbf{P}^{n_i}, \mathbf{P}^{n_{i+1}-1}, \dots, \mathbf{P}^{n_{j-1}-1}, \mathbf{P}^{n_j}, \mathbf{P}^{n_{j+1}-1}, \dots\}$

which have $\kappa \geq 1$ generators in common. Then, $d = d_1 + d_2$ and $g = g_1 + g_2 + \kappa - 1$.

Moreover, if $m = 0$, then the incidence scroll breaks up into a plane and an incidence scroll $R_g^{d-1} \subset \mathbf{P}^{n-1}$ with base $\ddot{\mathcal{B}}$ in general position.

Proof. By assumption we have a scroll formed by the lines which pass through $\mathbf{P}^m$ and a scroll generated by the lines which intersect the base spaces and lie in $\mathbf{P}^{n-1}$. The three scrolls are represented by curves $C = \bigcap_{k=1}^r \Omega(\mathbf{P}^{n_k}, \mathbf{P}^n)$, $\dot{C} = \Omega(\mathbf{P}^m, \mathbf{P}^n) \cap (\bigcap_{k \neq i, j} \Omega(\mathbf{P}^{n_k}, \mathbf{P}^n))$ and $\ddot{C} = \Omega(\mathbf{P}^{n_i}, \mathbf{P}^{n-1}) \cap \Omega(\mathbf{P}^{n_j}, \mathbf{P}^{n-1}) \cap (\bigcap_{k \neq i, j} \Omega(\mathbf{P}^{n_k-1}, \mathbf{P}^{n-1}))$ because every one of them satisfies (IS) .

Write $A_i = \Omega(\mathbf{P}^{n_i}, \mathbf{P}^n)$, $i = 1, \dots, r$, where the $\mathbf{P}^{n_i}$'s are in general position and $B_j = \Omega(\mathbf{P}^{n_j}, \mathbf{P}^n)$, $j = 1, \dots, r$, where $\mathbf{P}^{n_1} \cap \mathbf{P}^{n_2} = \mathbf{P}^m$ and the other subspaces are in general position. Then there is an invertible linear transformation of $\mathbf{P}^N$ into itself which carries $G(1, n)$ into itself and A_i into B_i (see [6], Proposition 4). Consequently, we conclude that A_i and B_i , $i = 1, \dots, r$, define the same cohomology class. In this way we obtain what the set of lines which simultaneously intersect the subspaces of $\mathcal{B}$ can be continuously deformed into the union of two set of lines. A set formed by the lines which pass through a fixed point and lie in a plane and other set formed by the lines which simultaneously intersect the subspaces of $\check{\mathcal{B}}$. Since the various subvarieties in a continuous system are all assigned the same cohomology class, we find the following equality of Schubert cycles, $C = \dot{C} + \ddot{C} = d\Omega(0, 2)$. So, $d = d_1 + d_2$.

We have in $G(1, n)$ a short exact sequence:

$$0 \rightarrow \mathcal{O}_C \longrightarrow \mathcal{O}_{\dot{C}} \oplus \mathcal{O}_{\ddot{C}} \longrightarrow \mathcal{O}_{\dot{C} \cap \ddot{C}} \rightarrow 0.$$

Therefore $\mathcal{X}(C) = \mathcal{X}(\dot{C}) + \mathcal{X}(\ddot{C}) - \mathcal{X}(\dot{C} \cap \ddot{C})$ where $\mathcal{X}(C)$ is the Euler characteristic, whence $g(C) = g(\dot{C}) + g(\ddot{C}) + \kappa - 1$. ■

Remark 3.2 Let $C_g^d = \bigcap_{k=1}^r \Omega(\mathbf{P}^{n_k}, \mathbf{P}^n)$ be a curve in $G(1, n)$, which defines an incidence scroll in $\mathbf{P}^n$. If we want to obtain another curve $\dot{C}$ which is an intersection of Schubert varieties with the same genus, then $\ddot{C}$ must be a line, there is only one generator in common and $\deg(\dot{C}) = d - 1$.

Let $D_g^{d_k}$ be the directrix curve of $R_g^d \subset \mathbf{P}^n$ contained in $\mathbf{P}^{n_k}$ for $k \in \{1, \dots, r\}$. Then the directrix curve of $R_g^{d-1} \subset \mathbf{P}^{n-1}$ in the $\mathbf{P}^{n_k-1}$ has degree $d_k - 1$, for $k \neq i, j$. This is not so in the other cases because if $k = i, j$, then the directrix curve in $\mathbf{P}^{n_k}$ has degree d_k (an easy computation of Schubert cycles).

We shall refer to this particular degeneration as join $\mathbf{P}^{n_i}$ and $\mathbf{P}^{n_j}$ and to the inverse as separate $\mathbf{P}^{n_i}$ and $\mathbf{P}^{n_j}$.

Example 3.3 The following examples are a good illustration of join and separate $\mathbf{P}^{n_i}$ and $\mathbf{P}^{n_j}$.

- $\mathcal{B}_1 = \{2\mathbf{P}^{n-1}, 2\mathbf{P}^n, \mathbf{P}^{2n-3}\} \implies R_{n-2}^{4n-6} \subset \mathbf{P}^{2n-1}$.

If we suppose $\mathbf{P}^n \cup \mathbf{P}^{2n-3} = \mathbf{P}^{2n-2}$, i.e., $\mathbf{P}^n \cap \mathbf{P}^{2n-3} = \mathbf{P}^{n-1}$, then the scroll breaks up into $R_0^{2n-2} \subset \mathbf{P}^{2n-1}$ with base $\{3\mathbf{P}^{n-1}, \mathbf{P}^n\}$ and $R_0^{2n-4} \subset \mathbf{P}^{2n-3}$ with base $\{3\mathbf{P}^{n-2}, \mathbf{P}^{n-1}\}$ (we will see it in section 4) and $n - 1$ generators in common. It follows that $d = 2n - 2 + 2n - 4 = 4n - 6$ and $g = n - 2$.

- $\mathcal{B}_2 = \{\mathbf{P}^{n-2}, 2\mathbf{P}^{n-1}, \mathbf{P}^n, \mathbf{P}^{2n-4}\} \implies R_{n-3}^{4n-9} \subset \mathbf{P}^{2n-2}$; apply 3.2 to a $\mathbf{P}^{n-1}$ and a $\mathbf{P}^n$ in $\mathcal{B}_1$ and write n instead of $n - 1$.
- $\mathcal{B}_3 = \{\mathbf{P}^{n-3}, 3\mathbf{P}^{n-1}, \mathbf{P}^{2n-5}\} \implies R_{n-4}^{4n-12} \subset \mathbf{P}^{2n-3}$; apply 3.2 to $\mathbf{P}^{n-2}$ and $\mathbf{P}^n$ in $\mathcal{B}_2$ and write n instead of $n - 1$.
- $\mathcal{B}_4 = \{3\mathbf{P}^{n-1}, \mathbf{P}^{n+1}, \mathbf{P}^{2n-3}\} \implies R_{n-3}^{4n-8} \subset \mathbf{P}^{2n-1}$; apply 3.2 to $2\mathbf{P}^{n-1}$'s in $\mathcal{B}_2$.

We will use the existence of the previous incidence scrolls in section 5.

4. Incidence Rational Scrolls

Let $X = \mathbf{P}(\mathcal{E})$ be a ruled surface. We say that X is a rational ruled surface if $C \cong \mathbf{P}^1$. One sees immediately that $X_0 = \mathbf{P}^1 \times \mathbf{P}^1$ with its first projection is a rational ruled surface. Each rational ruled surface is isomorphic to the ruled surface obtained from X_0

by applying a finite number of elementary transform at $P_1, \dots, P_e \in X_0$, with a suitable number $e \geq 0$. We shall write the above expression as $X_e = \text{elem}_{(P_1, \dots, P_e)}(X_0)$. The integer e is an invariant of X_e . Moreover, for each $e \geq 0$ there is exactly one (up to isomorphism) rational ruled surface with invariant e , given by $\mathcal{E} = \mathcal{O}_{\mathbf{P}^1} \oplus \mathcal{O}_{\mathbf{P}^1}(-e)$, i.e., $X_e \cong \mathbf{P}(\mathcal{E})$.

Proposition 4.1 *Let $D \sim aC_o + bf$ be a divisor on X_e . Then:*

1. D is very ample $\iff D$ is ample $\iff a > 0$ and $b > ae$;
2. $|D|$ contains an irreducible nonsingular curve $\iff |D|$ contains an irreducible curve $\iff a = 0, b = 1$; or $a = 1, b = 0$; or $a > 0, b > ae$; or $e > 0, a > 0, b = ae$. ■

Let H be a very ample divisor on X_e and let Φ_H be the closed immersion defined by $|H|$. Then we know that $\Phi_H(X_e)$ is a scroll if and only if $H \equiv C_o + mf$ with $m > e$. To see a more complete theory of rational ruled surfaces we refer the reader to [3] and [7].

The aim of this chapter is to study all the incidence rational scrolls with base in general position. For this, we can now formulate a theorem as follows.

Theorem 4.2 *Let $H \sim C_o + mf$ be a very ample divisor on X_e and let $\Phi_H : X_e \rightarrow R_0^{2m-e} \subset \mathbf{P}^{2m-e+1}$ be the closed immersion defined by $|H|$. Then R_0^{2m-e} is an incidence scroll if and only if it satisfies one of the following conditions:*

1. $H \sim C_o + (e+1)f$ ($\Phi_H(C_o) = \mathbf{P}^1$);
2. $e = 0$ ($\Phi_H(C_o) = C_0^m \subset \mathbf{P}^m$);
3. $e = 1$ ($\Phi_H(C_o) = C_0^{m-1} \subset \mathbf{P}^{m-1}$).

We will divide its proof into a sequence of propositions.

Rational Scrolls with a Directrix Line

It is clear that an incidence scroll with $\mathbf{P}^1$ as base space is a rational scroll. The following proposition gives all rational normal scrolls with a line as directrix. The affirmation is not

true if normal is deleted from the hypothesis. For example, $R_0^3 \subset \mathbf{P}^3$, which has a double line, is not an incidence scroll if the base spaces are in general position.

Proposition 4.3 *In $\mathbf{P}^n$ the incidence scroll with base $\mathcal{B}_n = \{\mathbf{P}^1, (n-1)\mathbf{P}^{n-2}\}$ is the rational normal scroll of degree $n-1$ with a line as minimum directrix.*

Proof. We proceed by induction in n . The proposition is true in $\mathbf{P}^3$. In this case, the scroll is the quadric surface in $\mathbf{P}^3$.

Supposing the proposition true in $\mathbf{P}^n$, we prove it for $n+1$. We can add a hyperplane to the base without affecting the scroll. Separating $\mathbf{P}^1$ and $\mathbf{P}^{n-1}$, as in 3.2, we obtain a base $\mathcal{B}_{n+1} = \{\mathbf{P}^1, n\mathbf{P}^{n-1}\}$ which defines $R_0^n \subset \mathbf{P}^{n+1}$ as incidence scroll. ■

Corollary 4.4 *For every $e \geq 0$, there is an embedding of X_e as incidence rational scroll.*

Proof. Use the very ample divisor $H \sim C_o + (e+1)f$. ■

Rational Scrolls of General Type

X_e is said to be of general type if and only if $\mathcal{E} \cong \mathcal{O}_{\mathbf{P}^1} \oplus \mathcal{O}_{\mathbf{P}^1}$ or $\mathcal{E} \cong \mathcal{O}_{\mathbf{P}^1} \oplus \mathcal{O}_{\mathbf{P}^1}(-1)$. Moreover, if X_e is general of degree d , then the minimum degree of the directrix curves is equal to $\frac{d}{2}$ (respectively, $\frac{d-1}{2}$) for d even (respectively, odd). On X_e , the family of directrices of degree m has dimension $d_m = 2m - d + 1$ (see [2], pp. 89-90).

Proposition 4.5 *All general rational scrolls are incidence scrolls.*

Proof. e=0: We will see that in $\mathbf{P}^{2m+1}$ the incidence scroll generated by $\mathcal{B} = \{3\mathbf{P}^m, \mathbf{P}^{m+1}\}$ is in fact $R_0^{2m} \subset \mathbf{P}^{2m+1}$. This incidence scroll have directrix curves in each $\mathbf{P}^m$ of degree d_m and a directrix curve in $\mathbf{P}^{m+1}$ of degree d_{m+1} . Then:

1. d_{m+1} is the number of lines in $\mathbf{P}^{2m+1}$ which meet $4\mathbf{P}^m$'s. Call this number $z(m)$ i.e.,

$$d_{m+1} = z(m) = \Omega(m, 2m+1)^4.$$

2. if we take a hyperplane $\mathbf{P}^{2m} \subset \mathbf{P}^{2m+1}$ through the base $\mathbf{P}^{m+1}$, then we can calculate

d_m as the number of generators of the scroll which meet $\mathbf{P}^{2m}$ in points of the base $\mathbf{P}^m$.

To obtain this number we consider two cases: either the generators lie in $\mathbf{P}^{2m}$ or they do

not. In the former case, we have the lines in $\mathbf{P}^{2m-3}$ which meet $4\mathbf{P}^{m-2}$'s therein. The

number of these lines is $z(m-2)$. In the latter case, this number is equal to the number

of lines through $\mathbf{P}^m \cap \mathbf{P}^{m+1}$, which meet the other $2\mathbf{P}^m$'s. Thus $d_m = z(m-2) + 1$.

Therefore $\deg(R) = d_m + d_{m+1} - 1 = 2d_m \implies z(m) = z(m-2) + 2 \implies d_{m+1} = z(m) = d_m + 1$. But $z(m)$ is the number of lines meeting $4\mathbf{P}^m$'s in $\mathbf{P}^{2m+1}$, i.e., by Schubert calculus, $z(m) = m + 1$. Then $\deg(R) = 2m$. Moreover, a scroll of degree $2m$ in $\mathbf{P}^{2m}$ is necessarily rational and normal.

e=1: Since $R_0^{2m-1} \subset \mathbf{P}^{2m}$ is of general type, it has one minimum directrix curve. Then we can take as base space a $\mathbf{P}^{m-1}$ which contains it. Choose $3\mathbf{P}^m$'s containing 3 generic directrix curves C_0^{m+1} . This is possible because $h^0(\mathcal{O}_{X_1}(C_o + f)) = 3$. By Proposition 4.1 and Bertini's Theorem we know that $|C_o + f|$ has irreducible curves and that the generic is irreducible. $\mathcal{B} = \{\mathbf{P}^{m-1}, 3\mathbf{P}^m\}$ is base of $R_0^{2m-1} \subset \mathbf{P}^{2m}$. We conclude it from 3.2, joining $2\mathbf{P}^m$ of case $e = 0$. ■

Rational Scrolls which are not Incidence Scrolls

Proposition 4.6 *Let $H \sim C_o + mf$ be a very ample divisor on X_e with $m > e + 1 > 2$. If the base must be in general position, then $\Phi_H(X_e) = R_0^{2m-e} \subset \mathbf{P}^{2m-e+1}$ cannot be obtained by incidences.*

Proof. Under the above hypotheses, we have $m + (e + 2)(m - e) > 4m - 2e - 1$ ($\iff 1 < e < m - 1$). Then:

$$m + (\eta)(m - e) \leq 4m - 2e - 1 \quad (*)$$

$$m + (\eta + 1)(m - e) > 4m - 2e - 1$$

for a suitable number η such that $1 < \eta < e + 2$. An argument similar to Proposition 2.7 shows that there are $\eta \mathbf{P}^m$'s in the base. Then $\mathcal{B} = \{\mathbf{P}^{m-e}, \eta \mathbf{P}^m, \mathbf{P}^{n_1}, \dots, \mathbf{P}^{n_r}\}$ with $m+1 \leq n_i \leq 2m-e-1$ is base of an incidence scroll, which is not rational. In other case, joining base spaces, we find a rational scroll in $\mathbf{P}^3$, obtained by Proposition 4.3, which is impossible. If we apply it successively the inverse degeneration (separate $\mathbf{P}^{n_i}$ and $\mathbf{P}^{n_j}$) to all the possible pairs (n_i, n_j) such that $\mathbf{P}^{n_i} \cap \mathbf{P}^{n_j} = \emptyset$, then we must obtain our scroll, but we obtain the rational normal scroll $R_0^{2m-e} \subset \mathbf{P}^{2m-e+1}$ with a line as minimum directrix curve (i.e., $m = e + 1$) or the rational scroll of general type (i.e., $e \leq 1$). Note that we can project to any $\mathbf{P}^{2m-e-k}$, and so join two base spaces, because the scroll is rational and, in particular, decomposable, i.e., $\mathbf{P}^{n_1} \cap \mathbf{P}^{n_2} = \emptyset$. If (*) is an equality, we apply this argument again, with $\mathcal{B}$ replaced by $\{\mathbf{P}^{m-e}, \eta \mathbf{P}^m\}$. Then no the rational scroll has a base which defines it as incidence scroll. ■

Corollary 4.7 *Let $R = R_0^{2m-e} \subset \mathbf{P}^{2m-e+1}$ be the projective model of X_e defined by $H \sim C_o + mf$. For $e \geq 1$, R is an incidence scroll (with base in general position) if and only if $mh^0(\mathcal{O}_X(C_o)) + (m-e)(h^0(\mathcal{O}_X(C_o + ef))) = 2(2m-e+1) - 3$.* ■

Example 4.8 Consider $R_0^6 \subset \mathbf{P}^7$ with a directrix conic, a three-dimensional family of directrix quartics and a five-dimensional one of directrix quintics. This surface is a projective model of X_2 by $H \sim C_o + 4f$. If we apply Corollary 4.7, then we see that it is not an incidence scroll because $4 + 4(4-2) \neq 2 \cdot 7 - 3$.

From Propositions 4.3, 4.5 and 4.6, we obtain the proof of Theorem 4.2. In these proofs, we have built a base for each incidence rational scroll in $\mathbf{P}^n$. It is easy notice that if a rational scroll is an incidence scroll, then there is really only one way to choose a base, i.e., the sequence $n_1, \dots, n_r$ is unique. Table 1 contains all the incidence rational scrolls up $\mathbf{P}^8$.

TABLE 1. INCIDENCE RATIONAL SCROLLS									
Scroll	$n_i, i = 1, \dots, 6$						Min. Dir. (*)	Normalized	$\deg(\mathfrak{b})$
	1	2	3	4	5	6			
$R_0^2 \subset \mathbf{P}^3$	3	-	-	-	-	-	$\mathbf{P}^1(\infty^1)$	$\mathcal{O}_{\mathbf{P}^1} \oplus \mathcal{O}_{\mathbf{P}^1}$	1
$R_0^3 \subset \mathbf{P}^4$	1	3	-	-	-	-	$\mathbf{P}^1(1)$	$\mathcal{O}_{\mathbf{P}^1} \oplus \mathcal{O}_{\mathbf{P}^1}(-1)$	2
$R_0^4 \subset \mathbf{P}^5$	1	-	4	-	-	-	$\mathbf{P}^1(1)$	$\mathcal{O}_{\mathbf{P}^1} \oplus \mathcal{O}_{\mathbf{P}^1}(-2)$	3
$R_0^4 \subset \mathbf{P}^5$	-	3	1	-	-	-	$C_0^2 \subset \mathbf{P}^2(\infty^1)$	$\mathcal{O}_{\mathbf{P}^1} \oplus \mathcal{O}_{\mathbf{P}^1}$	2
$R_0^5 \subset \mathbf{P}^6$	1	-	-	5	-	-	$\mathbf{P}^1(1)$	$\mathcal{O}_{\mathbf{P}^1} \oplus \mathcal{O}_{\mathbf{P}^1}(-3)$	4
$R_0^5 \subset \mathbf{P}^6$	-	1	3	-	-	-	$C_0^2 \subset \mathbf{P}^2(1)$	$\mathcal{O}_{\mathbf{P}^1} \oplus \mathcal{O}_{\mathbf{P}^1}(-1)$	3
$R_0^6 \subset \mathbf{P}^7$	1	-	-	-	6	-	$\mathbf{P}^1(1)$	$\mathcal{O}_{\mathbf{P}^1} \oplus \mathcal{O}_{\mathbf{P}^1}(-4)$	5
$R_0^6 \subset \mathbf{P}^7$	-	-	3	1	-	-	$C_0^3 \subset \mathbf{P}^3(\infty^1)$	$\mathcal{O}_{\mathbf{P}^1} \oplus \mathcal{O}_{\mathbf{P}^1}$	3
$R_0^7 \subset \mathbf{P}^8$	1	-	-	-	-	7	$\mathbf{P}^1(1)$	$\mathcal{O}_{\mathbf{P}^1} \oplus \mathcal{O}_{\mathbf{P}^1}(-5)$	6
$R_0^7 \subset \mathbf{P}^8$	-	-	1	3	-	-	$C_0^3 \subset \mathbf{P}^3(1)$	$\mathcal{O}_{\mathbf{P}^1} \oplus \mathcal{O}_{\mathbf{P}^1}(-1)$	4

* Number of minimum directrix curves

5. Incidence Elliptic Scrolls

Let $X = \mathbf{P}(\mathcal{E})$ be a ruled surface over an elliptic curve C . Let $\mathfrak{d}$ be a divisor on C . Then:

1. $\deg(\mathfrak{d}) \geq e + 2 \iff$ there is a section $D \sim C_o + \mathfrak{d}f$ and $|D|$ has no base points;
2. $\deg(\mathfrak{d}) \geq e + 3 \iff |C_o + \mathfrak{d}f|$ is very ample.

If we take $H \sim C_o + \mathfrak{b}f$ a very ample divisor on X with $m = \deg(\mathfrak{b})$, then we obtain the closed immersion $\Phi_H: X \longrightarrow R_1^{2m-e} \subset \mathbf{P}^{2m-e-1}$. We denote the scroll briefly by R .

We want to know if such a scroll is an incidence scroll. For this we need a base $\mathcal{B}$. Since such a base is formed by linear spaces which contain directrix curves of the scroll,

we begin to study the families of directrix curves in R . The aim of this section is to prove the following theorem.

Theorem 5.1 *Let $H \sim C_o + \mathfrak{b}f$ be a very ample divisor on X with $m = \deg(\mathfrak{b})$ and let $\Phi_H : X \rightarrow R_1^{2m-e} \subset \mathbf{P}^{2m-e-1}$ be the closed immersion defined by $|H|$. Then R_1^{2m-e} is an incidence scroll (with base in general position) if and only if it satisfies one of the following conditions:*

1. $e = -1$ and $m = 2$ ($\mathcal{B} = \{5\mathbf{P}^2\}$);
2. $\mathfrak{e} \sim 0$ and $m = 4$ ($\mathcal{B} = \{3\mathbf{P}^3, 2\mathbf{P}^5\}$);
3. X decomposable, $0 \leq e \leq 3$ and $m = e + 3$ ($\mathcal{B} = \{\mathbf{P}^2, (e+1)\mathbf{P}^{e+2}, (3-e)\mathbf{P}^{e+3}\}$).

Decomposable incidence elliptic scrolls

Let $X = \mathbf{P}(\mathcal{E})$ be a ruled surface over an elliptic curve C , defined by a normalized bundle $\mathcal{E} \cong \mathcal{O}_C \oplus \mathcal{O}_C(\mathfrak{e})$ and let $H \sim C_o + \mathfrak{b}f$ be a very ample divisor on X . Suppose R an incidence scroll with base in general position.

Consider $e \geq 1$. Since $h^0(\mathcal{O}_X(C_o)) = 1$, there does not exist another directrix curve of minimum degree. The following directrix curves of the scroll are linearly equivalent to $C_o - \mathfrak{e}f$ and, by H , are directrix curves of degree m . Since $h^0(\mathcal{O}_X(C_o - \mathfrak{e}f)) = e + 1 \geq 2$, we know that there are $(e+1)\mathbf{P}^{m-1}$'s in $\mathcal{B}$, whenever possible ($m-1+(e+1)(m-e-1) \leq 4m-2e-5$).

Lemma 5.2 *Let $X = \mathbf{P}(\mathcal{O}_C \oplus \mathcal{O}_C(\mathfrak{e}))$ be a ruled surface over an elliptic curve C with $e \geq 1$ and let $H \sim C_o + \mathfrak{b}f$ be a very ample divisor on X . Then R is an incidence scroll if and only if $\deg(\mathfrak{b}) = e + 3$ and $1 \leq e \leq 3$. Moreover, $\mathcal{B} = \{\mathbf{P}^2, (e+1)\mathbf{P}^{e+2}, (e-3)\mathbf{P}^{e+3}\}$.*

Proof. We know that there are $\mathbf{P}^{m-e-1}$ and $(e+1)\mathbf{P}^{m-1}$'s in $\mathcal{B}$ when $e = 1, 2, 3$. For $e = 1$, the following directrix curves are divisors $C \sim C_o + (P+Q)f$, $P, Q \in C$. Since

$h^0(\mathcal{O}_X(C_o + (Q - \mathfrak{e})f)) = 3$, we show that the generic curve is irreducible. For $m \geq 5$, we can see that one $\mathbf{P}^{m-2}$, $2\mathbf{P}^{m-1}$'s and one $\mathbf{P}^m$ are in $\mathcal{B}$, i.e., we impose $4m - 8$ independent conditions on $G(1, 2m - 2)$ so that $\mathcal{B} = \{\mathbf{P}^{m-2}, 2\mathbf{P}^{m-1}, \mathbf{P}^m, \mathbf{P}^{2m-4}\}$ is base of an incidence scroll. Such a scroll is of type $R_{m-3}^{4m-9} \subset \mathbf{P}^{2m-2}$ (see Example 3.3) which is not elliptic. For $m = 4$, $\mathcal{B} = \{\mathbf{P}^2, 2\mathbf{P}^3, 2\mathbf{P}^4\}$ is the base of $R_1^7 \subset \mathbf{P}^6$.

For $e = 2$, we only need another condition, i.e., $\mathcal{B} = \{\mathbf{P}^{m-3}, 3\mathbf{P}^{m-1}, \mathbf{P}^{2m-5}\}$. Such a scroll is $R_{m-4}^{4m-12} \subset \mathbf{P}^{2m-3}$ (see Example 3.3), which is elliptic if and only if $m = 5$.

For $e = 3$, we have seen that if R is an incidence scroll, then a $\mathbf{P}^{m-4}$ and $4\mathbf{P}^{m-1}$'s are in $\mathcal{B}$. If $m = 6$, $\mathcal{B} = \{\mathbf{P}^2, 4\mathbf{P}^5\}$ is base of $R_1^9 \subset \mathbf{P}^8$. For $e \geq 3$ and $m \neq 6$, we have $m - 1 + (e + 1)(m - e - 1) > 4m - 2e - 5$. Consequently, we cannot take $(e + 1)\mathbf{P}^{m-1}$'s as base spaces because these impose too many conditions on $G(1, 2m - e - 1)$. Since $m - 1 < 2m - e - 3$, we conclude (as in Proposition 4.6) that R is not an incidence scroll. ■

Remark 5.3 We have seen that, in $\mathbf{P}^N$, if the base space of greater dimension is $\mathbf{P}^{N-2}$'s, then it is not necessary to take the number of linearly independent directrix curves. From this, we need to show if $m - 1 < 2m - e - 3$.

We now evaluate the case $e = 0$. There are two choices of the divisor $\mathfrak{e}$. If $\mathfrak{e} \not\sim 0$, there are two choices of normalized $\mathcal{E}$, namely $\mathcal{O}_C \oplus \mathcal{O}_C(\mathfrak{e})$ and $\mathcal{O}_C \oplus \mathcal{O}_C(-\mathfrak{e})$. There are exactly two choices of C_o , both with $C_o^2 = 0$, namely C_o and $C_o - \mathfrak{e}f$. Since $h^0(\mathcal{O}_X(C_o)) = h^0(\mathcal{O}_C(C_o - \mathfrak{e}f)) = 1$, these are the only minimum directrix curves. If R is an incidence scroll with base in general position, then there are $2\mathbf{P}^{m-1}$'s in $\mathcal{B}$. The following directrix curves are in linear systems $|C_o + (P + \mathfrak{e})f|$ and $|C_o + (P - \mathfrak{e})f|$, $P \in C$, where $h^0(\mathcal{O}_X(C_o + (P + \mathfrak{e})f)) = h^0(\mathcal{O}_X(C_o + (P - \mathfrak{e})f)) = 2$. As in the proofs described above we can see that, if R is an incidence scroll, there are $2\mathbf{P}^m$'s as base spaces. Then $\mathcal{B} = \{2\mathbf{P}^{m-1}, 2\mathbf{P}^m, \mathbf{P}^{2m-3}\}$ and the incidence scroll has degree $4m - 6$ and genus $m - 2$ (see Example 3.3), which is elliptic if and only if $m = 3$.

In the other case, $\mathfrak{e} \sim 0$, there is a one-dimensional family of directrix curves of degree m . If R is an incidence scroll, there are $3\mathbf{P}^{m-1}$'s in $\mathcal{B}$. The following directrix curves are in a linear system $|C_o + (P+Q)f|$ where $h^0(\mathcal{O}_X(C_o + (P+Q)f)) = 4$. By Bertini, we know that it has irreducible curves and that so is the generic curve. If R is an incidence scroll, then $m \geq 4$ (no three $\mathbf{P}^{m-1}$'s suffice and another $\mathbf{P}^{m-1}$ imposes too many conditions). For $m \geq 4$, $\mathcal{B} = \{3\mathbf{P}^{m-1}, \mathbf{P}^{m+1}, \mathbf{P}^{2m-3}\}$ generates a scroll of genus $m-3$ (see Example 3.3), which is elliptic if and only if $m = 4$.

Remark 5.4 If we want to extend the study of incidence elliptic scrolls to incidence scrolls of genus g , then we must consider the following properties.

1. If the scroll has a unique minimum directrix curve, then the space which contains it is in $\mathcal{B}$.
2. If the scroll has two minimum directrix curves, then the correspondent spaces are in $\mathcal{B}$.
3. If the scroll has an one-dimensional family of minimum directrix curves, then the base has three spaces which contain three of these curves.
4. If $m - g + (e + 2 - g)(m - e - g) > 4m - 2e - 4g - 1$, then R is not an incidence scroll ($g = 0, 1$).

Indecomposable incidence elliptic scrolls

Let $X = \mathbf{P}(\mathcal{E})$ be a ruled surface over an elliptic curve C , corresponding to an indecomposable $\mathcal{E}$. Then $e = 0, -1$ and there is exactly one ruled surface over C for each of these values of e . Let $H \sim C_o + \mathfrak{b}f$ be a divisor with $\deg(\mathfrak{b}) = m \geq e + 3$, which defines $\Phi_H: X \longrightarrow R_1^{2m-e} \subset \mathbf{P}^{2m-e-1}$. We want to know if such a scroll is an incidence scroll. For this, we need to find a suitable number of base spaces in $\mathbf{P}^{2m-e-1}$ whose dimension is $\leq 2m - e - 3$

If $e = 0$ then $\mathcal{E}$ is an extension of $\mathcal{O}_C$ by $\mathcal{O}_C$ i.e. given by an exact sequence $0 \rightarrow \mathcal{O}_C \rightarrow \mathcal{E} \rightarrow \mathcal{O}_C \rightarrow 0$. Since $C_0^2 = 0$ and X is indecomposable, there is a unique directrix of minimum degree equal to m . For any $P \in C$, $h^0(\mathcal{O}_X(C_o + Pf)) = 2$. Then we have an one-dimensional family $|C_o + Pf|$ of directrix curves of degree $m + 1$, for any $P \in C$. Suppose that R is an incidence scroll with base in general position. An easy computation shows that $\mathcal{B} = \{\mathbf{P}^{m-1}, 2\mathbf{P}^m, \mathbf{P}^{n_1}, \dots, \mathbf{P}^{n_r}\}$ where $m \leq n_i \leq 2m - 3$ and $\sum_{i=1}^r (2m - 2 - n_i) = m$. Then we can obtain another incidence elliptic scroll in $\mathbf{P}^{2m}$ with $e = 1$ and base $\dot{\mathcal{B}} = \{\mathbf{P}^{m-1}, \mathbf{P}^m, \mathbf{P}^{m+1}, \mathbf{P}^{n_1+1}, \dots, \mathbf{P}^{n_r+1}\}$ (separate $\mathbf{P}^{m-1}$ and $\mathbf{P}^m$ in $\mathcal{B}$). This happens when $m = 3$, $n_1 = 2$ and $n_2 = 3$, which contradicts the hypothesis $m \leq n_i$. Hence there are no indecomposable elliptic scrolls with $e = 0$ defined by incidences.

If $e = -1$ then we have an exact sequence $0 \rightarrow \mathcal{O}_C \rightarrow \mathcal{E} \rightarrow \mathcal{O}_C(P) \rightarrow 0$. Then $\Phi_H(C_o)$ is a directrix curve of degree $m + 1$. If D is a section with $D^2 = 1$, then $D \sim C_o + (\delta - \mathfrak{e})f$ with $\deg(\delta) = 1$. In fact, the sections C_o with $C_o^2 = 1$ form a one-dimensional algebraic family parametrized by C such that no two of them are linearly equivalent. If R is an incidence scroll, then $\mathcal{B} = \{2\mathbf{P}^m, \mathbf{P}^{n_1}, \dots, \mathbf{P}^{n_r}\}$ with $m \leq n_i \leq 2m - 2$ and $\sum_{i=1}^r (2m - n_i - 1) = 2m - 1$. We can obtain another incidence elliptic scroll in $\mathbf{P}^{2m+1}$ with $\mathfrak{e} \neq 0$ and $e = 0$ (separate $2\mathbf{P}^m$'s in $\mathcal{B}$). Consequently, there is only one incidence scroll with $e = -1$. This happens when $m = n_i = 2$ and $r = 3$. So $\Phi_H(X) = R_1^5 \subset \mathbf{P}^4$ is an incidence scroll with base $\mathcal{B} = \{5\mathbf{P}^2\}$.

In Table 2 we have compiled all incidence elliptic scrolls. The interest of this table is that each scroll is a degenerate scroll of the remainder of them.

TABLE 2. INCIDENCE ELLIPTIC SCROLLS							
Scroll	$n_i, i = 1, \dots, 4$				Min. Dir.	Normalized	$\deg(\mathbf{b})$
	2	3	4	5			
$R_1^5 \subset \mathbf{P}^4$	5	-	-	-	$C_1^3 \subset \mathbf{P}^2$	$\mathcal{E} \in \text{Ext}^1(\mathcal{O}_C(P), \mathcal{O}_C)$	2
$R_1^6 \subset \mathbf{P}^5$	2	3	-	-	$C_1^3 \subset \mathbf{P}^2$	$\mathcal{O}_C \oplus \mathcal{O}_C(\mathbf{e})(\mathbf{e} \neq 0)$	3
$R_1^7 \subset \mathbf{P}^6$	1	2	2	-	$C_1^3 \subset \mathbf{P}^2$	$\mathcal{O}_C \oplus \mathcal{O}_C(-P)$	4
$R_1^8 \subset \mathbf{P}^7$	1	-	3	1	$C_1^3 \subset \mathbf{P}^2$	$\mathcal{O}_C \oplus \mathcal{O}_C(-P-Q)$	5
$R_1^8 \subset \mathbf{P}^7$	-	3	-	2	$C_1^4 \subset \mathbf{P}^3$	$\mathcal{O}_C \oplus \mathcal{O}_C$	4
$R_1^9 \subset \mathbf{P}^8$	1	-	-	4	$C_1^3 \subset \mathbf{P}^2$	$\mathcal{O}_C \oplus \mathcal{O}_C(-P-Q-R)$	6

Acknowledgements: Rosa Cid-Muñoz was supported by a fellowship of Xunta de Galicia (Spain).

References

- [1] EDGE, W. L. *The Theory of Ruled Surfaces*.
Cambridge Univesity Press. (1931).
- [2] GHIONE, F. *Quelques rèsultats de Corrado Segre sur les sufaces reglees*.
Math. Ann. 255, pp. 77-95 (1981).
- [3] HARTSHORNE, R. *Algebraic Geometry*.
Graduate Texts in Mathematics 52. Springer-Verlag, New York. 1977.
- [4] IITAKA, S. *Algebraic Geometry. An Introduction to Birational Geometry of Algebraic Varieties*.
Graduate Texts in Mathematics 76. Springer-Verlag, New York Heidelberg Berlin. 1982.
- [5] HERNÁNDEZ, R. - SOLS, I. *Connectedness of intersections of special Schubert varieties*.
Manuscripta Mathematica 83, pp. 215-222 (1994).
- [6] KLEIMAN, S. L. - LAKSOV, D. *Schubert calculus*.
Amer. Math. Monthly 79, pp. 1061-1082 (1972).
- [7] LANTERI, A. -PALLESCHI, M. *Sulle rigate geometrichi razionali ed alcune questioni di immersione*.
Milano, Istituto matematico "F. Enriques". Quaderno n.13/s. (1978).